

STUDYING THE CLINICAL AND IMMUNOLOGICAL NATURE OF THE COURSE OF FORMS OF SYPHILIS

Oripov Rustam Anvarovich

Samarkand State Medical University Department of Skin and Venereal

Abstract: *Enzyme immunoassay (ELISA) is one of the modern serological tests for syphilis and is used to diagnose both current and past syphilitic infection. The determination of antitreponemal IgG and IgM is of crucial importance for the diagnosis of early forms of congenital syphilis, differential diagnosis of reinfection-relapses, as well as for the study of the causes of seroresistance.*

Keywords: *Immunoassay, syphilis.*

Introduction: Syphilis is characterized by a fairly high number of patients, medical and social significance, difficulties in diagnosis and treatment. In the 90s, the fifth "wave" of the syphilis epidemic began in Russia, coinciding with unfavorable socio-economic factors, the growth of prostitution, and the abolition of legislative acts on the control and prevention of sexually transmitted infections. In 1991, the incidence of syphilis in the country per 100 thousand population was 7.2, and in 1997 it was 277.3, i.e. it increased 38.5 times, reaching its peak! In subsequent years, there was a gradual decrease in the incidence, however, and in 2013 the indicator is still high - 28.9. In 1997-2013, the incidence of syphilis in the country is 4 times higher than the initial 1991. It is natural that during the period of an increase in the incidence of syphilis, its early forms – Lues I and Lues II recens (or as it is customary to say today – secondary syphilis with a prescription of up to 6 months) prevail, with a decrease – hidden as early - with a prescription of up to 2 years, and later - more than 2 years. Latent forms of infection (both early and especially late) are insidious because they are a reservoir of possible neurosyphilis, its visceral forms. The second reservoir of these

crippling forms of syphilis are patients with seroresistance. Currently, seroresistance is understood as the state of the body, which is characterized by the absence of a decrease in the titers of reagins in the RMP (microprecipitation reaction) by 4 times or more within a year after the end of full-fledged specific treatment of syphilis. What has contributed to the increase in such a contingent of people since 1995? The epidemic of syphilis in the country required the transition to outpatient treatment of patients with this infection. Due to the lack of own research, registration of foreign benzatin-benzylpenicillin (extensillin, retarpen) was carried out, which allowed this transition to be made after the publication of the "Information Letter" in 1995. This official document allowed all infectious forms of syphilis (Lues I, Lues II, Lues latens praecox) to be treated with these reparations, including pregnant women. Of course, treatment with benzatin-benzylpenicillin has played a positive role in the fight against the epidemic. However, in the future, cases of seroresistance began to appear and most often in patients with Lues II recidiva and with Lues latens praecox. In our opinion, back in that distant 1998, the treatment of the above-mentioned forms of syphilis with durant drugs should have been banned in Russia. Nowadays, it should be remembered that seroresistance is dangerous in its consequences. When performing a spinal puncture in this category of persons, changes in the cerebrospinal fluid (positive serological reactions, increased protein content, cytolysis), combined with neurological symptoms, make it possible to establish neurosyphilis in about half of the patients.

The aim of the study was to determine the frequency of detection of antitreponemal IgG and Ig M in seroresistant syphilis.

Material and methods of research: The blood sera of 25 patients (14 men and 11 women) were examined, in whom serological reactions remained positive after the end of treatment. The main age category was a group of patients aged 20 to 40 years - 12 people, 8 patients aged 40 to 50 years, 5 patients aged 50 to 60 years. The ELISA procedure was performed according to the instructions for use attached to the test system (TS), the optical density of the samples was recorded using a flatbed spectrophotometer (Multiscan, Finland).

In terms of reaction mechanism, sensitivity and specificity, ELISA is close to immunofluorescence (RIF) reactions. The participation of the same antibodies in these reactions, a close titer level, and a high frequency of matching results were noted. Therefore, the results of Ig M-ELISA were compared with DAC and RIBT

Results: To detect IdM in people with seroresistance, it is advisable to use test systems characterized by high sensitivity in the late stages of syphilis. It is not recommended to use test systems designed only for early diagnosis of syphilis to detect IdM in people with seroresistance. Since according to the literature, the sensitivity of such test systems in seroresistant syphilis is about 6%. The results of ELISA in the group of patients with seroresistant syphilis are in good agreement with the ideas of suppressing the expression of IdM after adequate specific therapy while maintaining the synthesis and circulation of Ig G in the blood.

It should be noted that all 25 sera were IgM negative, but IgG- positive with varying degrees of optical density (OP). 18 samples had OP above 1.0 and 7 samples had OP below 1.0 Traditional serological tests were positive.

Conclusion: One of the main measures aimed at preventing an increase in the incidence of syphilis is the early diagnosis of this infection. Due to the development of more sensitive, specific and less time-consuming serological methods, the use of ELISA has become possible, as well as its use for confirmation in patients with seroresistance in syphilis.

LITERATURE

1. Islamov N. H., Yusupov M. M., Oripov R. A. CHEKLANGAN SKLERODERMANI DAVOLASHNING SAMARALI //Educational Research in Universal Sciences. – 2023. – T. 2. – №. 14. – C. 324-329.
2. Mardonovich N. R. et al. Epidemiological Analysis of Trichofitia Disease in Samarkand Region //Journal of Survey in Fisheries Sciences. – 2023. – T. 10. – №. 1S. – C. 4981-4994.

3. Anvarovich O. R., Ravshanovich K. K. A New View on the Course of Sexually Transmitted Diseases //Web of Semantic: Universal Journal on Innovative Education. – 2023. – Т. 2. – №. 2. – С. 189-193.
4. Anvarovich O. R., Javlonovna P. A. Takrakor in the Treatment of Itching Dermatitis //Web of Semantic: Universal Journal on Innovative Education. – 2023. – Т. 2. – №. 2. – С. 184-188.
5. Anvarovich O. R. Management of Sick Children with Adopic Dermatitis //Web of Semantic: Universal Journal on Innovative Education. – 2023. – Т. 2. – №. 2. – С. 179-183.
6. Oripov R. A. LABORATORY PARAMETERS IN CHILDREN WITH VERIFIED MENINGITIS //Modern Scientific Research International Scientific Journal. – 2023. – Т. 1. – №. 7. – С. 162-168.
7. Davlyatovna K. G., Anvarovich O. R., Xikmatovich I. N. UROGENITAL XLAMIDIYADA MIRAMISTINNING SAMARADORLIGINI BAHOLASH //Нововведения Современного Научного Развития в Эпоху Глобализации: Проблемы и Решения. – 2023. – Т. 1. – №. 4. – С. 53-59.
8. Kurbonalievich A. S. et al. UROGENITAL CHLAMYDIOSIS IS A CAUSE OF STERILITY AND CHRONIC PROSTATITIS IN MEN //Web of Scholars: Multidimensional Research Journal. – 2022. – Т. 1. – №. 6. – С. 306-310.
9. Anvarovich O. R. et al. ETIOPATHOGENESIS AND CLINICAL COURSE SPECIFIC CHARACTERISTICS OF NEURODERMITIS DISEASE //Web of Scholars: Multidimensional Research Journal. – 2022. – Т. 1. – №. 6. – С. 276-280.
10. Bakhodirovich T. I. et al. COMBINED IMMUNOTHERAPY IN THE TREATMENT OF CANDIDIASIAN INFECTIONS OF THE URINARY ORGANS //Web of Scholars: Multidimensional Research Journal. – 2022. – Т. 1. – №. 6. – С. 257-263.
11. Anvarovich O. R. ATOPIK DERMATIT BILAN BIRGA KECHADIGAN DIABET KASALLIGINI DAVOLASH SAMARADORLIGINI BAHOLASH

//BARQARORLIK VA YETAKCHI TADQIQOTLAR ONLAYN ILMIIY JURNALI.
– 2022. – С. 242-244.

12. Anvarovich O. R. APPLICATION OF VIFIRON FOR LOCAL TREATMENT OF HERPIS VIRUS //Евразийский журнал медицинских и естественных наук. – 2022. – Т. 2. – №. 3. – С. 30-32.

13. Kurbonalievich A. S. et al. Experience of the Combination of Tiflox and Immunomax in the Treatment of Trichomoniasis Combined with a Bacterial Process //Annals of the Romanian Society for Cell Biology. – 2021. – С. 2376-2380.

14. Kurbonalievich A. S. et al. Some Features of Psoriasis Therapy in Stationary //European Scholar Journal. – 2021. – Т. 2. – №. 3. – С. 248-250.

15. Kurbonalievich A. S. et al. Careful Attention To The History Of Chronic Urticaria Is One Of The Important Factors Of Productive Therapy //The American Journal of Medical Sciences and Pharmaceutical Research. – 2021. – Т. 3. – №. 02. – С. 55-58.

16. Kurbonalievich A. S. et al. Features Of The Use Of Antioxidant Drugs In The Treatment Of Psoriatic Disease //The American Journal of Medical Sciences and Pharmaceutical Research. – 2021. – Т. 3. – №. 02. – С. 50-54.

17. Орипов Р. А. и др. ОСОБЕННОСТИ ПРИМЕНЕНИЕ АНТИОКСИДАНТНЫХ ПРЕПАРАТОВ В ЛЕЧЕНИИ НЕЙРОДЕРМИТА. – 2021.

18. Ахмедов Ш. К. и др. НЕКОТОРЫЕ ОСОБЕННОСТИ ТАКТИКИ ПРОДУКТИВНОГО ЛЕЧЕНИЯ ХРОНИЧЕСКОЙ КРАПИВНИЦЫ. – 2021.

19. Salamova L. A. et al. Cutaneous leishmaniasis and concomitant tissue helminthiasis //ResearchJet Journal of Analysis and Inventions. – 2021. – Т. 2. – №. 03. – С. 64-72.

20. Ачилова О. Д. и др. Эхинококкоз касаллигини иммуноташхислашда антигенларни қўллашнинг тамоиллари //Общество и инновации. – 2020. – Т. 1. – №. 1/s. – С. 598-603.

21. Tolibov M. M., qizi Buronova F. Z. MAHALLIY OZON TERAPIYASINI QO‘LLAGAN HOLDA KOMPLEKS DAVOLASH FONIDA YOSH ERKAKLARDA SURUNKALI URETROGEN PROSTATIT (CUP) KECHISHINI O‘RGANISH //Educational Research in Universal Sciences. – 2024. – T. 3. – №. 1. – C. 417-421.
22. Tolibov M. M., Shodmonova M. A. BOLALARDA ATOPIK DERMATIT (AD) PATOGENEZIDA GENETIK OMILLARNING ROLINI BAHOLASH VA IMMUNOLOGIK PARAMETRLARGA QARAB ANIQLANGAN POLIMORFIZMLARNING PROGNOZIK AHAMIYATINI ANIQLASH //Educational Research in Universal Sciences. – 2024. – T. 3. – №. 1. – C. 406-410.
23. Tolibov M. M. et al. ATOPIK DERMATITNI DAVOLASHDA LAZER TERAPIYASINING SAMARADORLIGI //Educational Research in Universal Sciences. – 2024. – T. 3. – №. 1. – C. 286-289.
24. Utayev A. J., Tolibov M. M., Axmedova M. M. MANAGEMENT APPROACHES FOR CHRONIC URETHROGENIC PROSTATITIS //Modern Scientific Research International Scientific Journal. – 2024. – T. 2. – №. 1. – C. 151-156.
25. Makhmudovich T. M. ATOPIC DERMATITIS INCIDENT LEVEL, DIAGNOSTIC CRITERIA AND PROGNOSTIC IMPORTANCE //JOURNAL OF APPLIED MEDICAL SCIENCES. – 2023. – T. 6. – №. 4. – C. 158-168.
26. Толибов М. М. СОВРЕМЕННОЕ СОСТОЯНИЕ ПРОБЛЕМ ЭПИДЕМИОЛОГИИ, МИКРОБИОЛОГИИ И ТЕРАПИИ АКНЕ (ОБЗОР ЛИТЕРАТУРЫ) //SCIENTIFIC APPROACH TO THE MODERN EDUCATION SYSTEM. – 2023. – T. 2. – №. 14. – C. 215-224.
27. Ilyayevna P. U. et al. DERMATOSCOPIC PAINTING SOME COMMON DERMATOSES //World scientific research journal. – 2023. – T. 16. – №. 2. – C. 107-120.
28. Ilyayevna P. U. et al. MODERN TRENDS IN TREATMENT SICK MICROSPORIA //World scientific research journal. – 2023. – T. 16. – №. 2. – C. 98-106.

29. Kholikovich E. S. et al. APPLICATION OF" BETASALIK" OINTMENT IN THE THERAPY OF CHRONIC INFLAMMATORY AND ALLERGIC DERMATOSES //Journal of new century innovations. – 2023. – T. 31. – №. 3. – С. 56-62.